

Nutrazu® Flamingo Complete

50W3

Description

Nutrazu® Flamingo Complete is a complementary feedingstuff for all captive flamingos.

Features and Benefits

- Minimum wastage - Nutrazu® Flamingo Complete floats, thereby avoiding the accumulation of food on the bottom, which crustacean-eating birds have difficulty consuming.
- Nutritionally balanced - contains high levels of vitamins and fish proteins essential for growth and maintenance.
- Contains canthaxanthin - A pigment well dispersed within the feed that maintains plumage color and avoids the wastage of high-value pigment common with other methods of administration.

Product Form

- Extruded pellet size: $\frac{3}{16}$ " diameter x $\frac{3}{8}$ " length.. 20 kg. net weight paper sack.

Guaranteed Analysis

Crude protein not less than	19.0%
Crude fat not less than	5.0%
Crude fibre not more than	4.0%
Calcium (Ca) not less than	1.20%
Calcium (Ca) not more than	1.70%
Phosphorus (P) not less than	0.70%
Sodium (Na) not less than	0.10%
Sodium (Na) not more than	0.60%

Ingredients

Ground corn, wheat middlings, ground wheat, fish meal, porcine meat meal, dehulled soybean meal, brewers dried yeast, soybean oil, calcium carbonate, blood meal, dicalcium phosphate, monocalcium phosphate, DL-methionine, L-lysine, salt, pyridoxine hydrochloride, choline chloride, cholecalciferol (vitamin D₃), dl-alpha tocopheryl acetate (vitamin E), canthaxanthin, vitamin A acetate, menadione dimethylpyrimidinol bisulfite (vitamin K), L-ascorbyl-2-polyphosphate (vitamin C), biotin, ethoxyquin (a preservative), thiamin mononitrate, calcium pantothenate, riboflavin, magnesium oxide, nicotinic acid, folic acid, d-alpha tocopheryl acetate (natural source vitamin E), inositol, vitamin B₁₂ supplement, manganous oxide, p-aminobenzoic acid, zinc oxide, copper sulfate, calcium iodate, sodium selenite.

Feeding Directions

Introduce gradually until it forms the major part of the daily intake for growth and or maintenance. Feed at 23-30 g/kg body weight per bird per day. Increase feed to compensate for food taken by non-collection birds. Typical intake is 150-200g / day

(Continued on Back)

(Continued from Front)

Feed in shallow trough or in shallow water.

No need for additional colour supplementation.

For further information refer to www.Nutrazu.com.

Store in a cool dry place and fold down top when not in use

Approximate Nutrient Composition

NUTRIENTS

Protein, %	20.1
Arginine, %	1.10
Cystine, %	0.26
Glycine, %	1.20
Histidine, %	0.51
Isoleucine, %	0.74
Leucine, %	1.51
Lysine, %	1.26
Methionine, %	0.94
Phenylalanine, %	0.83
Tyrosine, %	0.52
Threonine, %	0.70
Tryptophan, %	0.22
Valine, %	0.94
Fat, %	5.5
Linoleic Acid, %	2.4
Fibre (Crude), %	3.6
Neutral Detergent Fibre, %	15.7
Acid Detergent Fibre, %	4.5
Metabolisable Energy, kcal/g	3.14

Quality Controlled by PMI Nutrition International, a subsidiary of America's oldest and largest animal nutrition company.

MINERALS

Ash, %	6.9
Calcium, %	1.47
Phosphorus, %	1.00
Phosphorus (non-phytate), %	0.69
Potassium, %	0.71
Magnesium, %	0.21
Sodium, %	0.16
Chlorine, %	0.31
Sulfur, %	0.36
Iron, ppm	280
Zinc, ppm	120
Manganese, ppm	140
Copper, ppm	13
Iodine, ppm	1.9
Selenium, ppm	0.30
Cobalt, ppm	0.28

VITAMINS

Vitamin K (as menadione), ppm	0.90
Thiamin Hydrochloride, ppm	19
Riboflavin, ppm	11
Niacin, ppm	110
Pantothenic Acid, ppm	24
Choline Chloride, ppm	1,580
Folic Acid, ppm	2.1
Pyridoxine, ppm	15
Biotin, ppm	0.40
Vitamin B ₁₂ , mcg/kg	24
Vitamin A, IU/gm	23
Vitamin D ₃ (added), IU/gm	4.6
Vitamin E, IU/kg	310
Canthaxanthin, ppm	40